

TEXAS CHILDREN'S HOSPITAL EVIDENCE-BASED OUTCOMES CENTER


High Flow Nasal Cannula (HFNC) Therapy: Initiation and Escalation for Respiratory Distress Evidence-Informed Pathway

Target Group
See TCH Bronchiolitis Guideline
Consider for other age groups and disease entities, such as asthma and CAP

Table 1: Clinical Respiratory Score (CRS)

Assess	Clinical Respiratory Score (CRS)		
	Score 0	Score 1	Score 2
Respiratory Rate	< 2 mos: < 50 2-12 mos: < 40 1-5 yrs: < 30 > 5 yrs: < 20	< 2 mos: 50-60 2-12 mos: 40-50 > 1-5 yrs: 30-40 > 5 yrs: 20-30	< 2 mos: > 60 2-12 mos: > 50 > 1-5 yrs: > 40 > 5 yrs: > 30
Auscultation	Good air movement, scattered expiratory wheezing, loose rales/crackles	Depressed air movement, inspiratory and expiratory wheezes or rales/crackles	Diminished or absent breath sounds, severe wheezing, or rales/crackles, or marked prolonged expiration
Use of Accessory Muscles	Mild to no use of accessory muscles, mild to no retractions, no nasal flaring on inspiration	Moderate intercostal retractions, mild to moderate use of accessory muscles, nasal flaring	Severe intercostal and subcostal retractions, nasal flaring
Mental Status	Normal to mildly irritable	Irritable, agitated, restless	Lethargic
Room Air SpO₂	> 95%	90-95%	< 90%
Color	Normal	Pale to normal	Cyanotic, dusky

(Add score from all rows to calculate total CRS)



****Patient disposition should NOT be based on HFNC settings (i.e., FiO₂, flow). Patient disposition should be determined by the overall clinical condition, which is mainly defined by CRS. See next page for additional guidance.**

Clinical standards are developed for 80% of the patient population with a particular disease. Each practitioner must use his/her clinical judgment in the management of any specific patient.

Critical Points of Evidence

Evidence Supports

- Use HFNC therapy in children experiencing respiratory distress. Use the maximum flow rate for the patient's appropriate cannula size. (1-12) – Strong recommendation, low quality evidence
- Identify nonresponders as patients exhibiting no response (e.g., HR, RR) within 1 hour of therapy. (13-16) – Strong recommendation, low quality evidence
The clinical respiratory score (CRS) used at TCH includes respiratory rate, among other markers. Patients with a significant cardiopulmonary disorder may have a higher HFNC therapy failure rate than the general population.

Evidence Lacking/Inconclusive

- Utilize the [Oxygen Weaning Protocol](#) for HFNC therapy weaning. – Consensus recommendation

Table 1: Exacerbation Severity Assessment Tool- Clinical Respiratory Score (CRS)

Clinical Respiratory Score (CRS)			
Assess	Score 0	Score 1	Score 2
Respiratory Rate	< 2 mos: < 50 2-12 mos: < 40 1-5 yrs: < 30 > 5 yrs: < 20	< 2 mos: 50-60 2-12 mos: 40-50 > 1-5 yrs: 30-40 > 5 yrs: 20-30	< 2 mos: > 60 2-12 mos: > 50 > 1-5 yrs: > 40 > 5 yrs: > 30
Auscultation	Good air movement, scattered expiratory wheezing, loose rales/crackles	Depressed air movement, inspiratory and expiratory wheezes or rales/crackles	Diminished or absent breath sounds, severe wheezing, or rales/crackles, or marked prolonged expiration
Use of Accessory Muscles	Mild to no use of accessory muscles, mild to no retractions, no nasal flaring on inspiration	Moderate intercostal retractions, mild to moderate use of accessory muscles, nasal flaring	Severe intercostal and substernal retractions, nasal flaring
Mental Status	Normal to mildly irritable	Irritable, agitated, restless.	Lethargic
Room Air SpO₂	> 95%	90-95%	< 90%
Color	Normal	Pale to normal	Cyanotic, dusky

(Add score from all rows to calculate total CRS)

Table 2: Inclusion/exclusion Criteria for Acute Care Areas

The following are general admission/exclusion criteria for acute care areas and are not exclusive to this protocol. These are provided to assist and offer **general guidance** on patient disposition and are **not** meant to be **all-inclusive**. **Patient needs and status** will ultimately **determine disposition** and will be based on discussion amongst the multidisciplinary providers (i.e., RT, physician, nurse).

Main Campus Acute Care	<p>Inclusion:</p> <ul style="list-style-type: none"> ○ CRS 0-5 at time of disposition and/or transfer, <i>if stable or improving on allowable max. therapies</i> <p>Exclusion:</p> <ul style="list-style-type: none"> ○ Patient <i>is not</i> stable or improving on allowable max. therapies ○ Patient requiring continuous albuterol therapy ○ CPAP or BiPAP use for patients with <i>acute</i> respiratory disease
MC Respiratory Unit	<p>Inclusion:</p> <ul style="list-style-type: none"> ○ Patient requiring continuous albuterol therapy ○ CRS 0-5 at time of disposition and/or transfer, <i>if stable or improving on allowable max. therapies</i> <p>Exclusion:</p> <ul style="list-style-type: none"> ○ CPAP or BiPAP use for patients with <i>acute</i> respiratory disease <p>Need for <i>additional</i> magnesium doses or terbutaline infusion</p>
WC Acute Care	<p>Inclusion:</p> <ul style="list-style-type: none"> ○ Patient requiring continuous albuterol therapy ○ CRS 0-5 at time of disposition and/or transfer, <i>if stable or improving on allowable max. therapies</i> <p>Exclusion:</p> <ul style="list-style-type: none"> ○ Patient <i>is not</i> stable or improving on allowable max. therapies ○ CPAP or BiPAP use for patients with <i>acute or chronic</i> respiratory disease
Woodlands Acute Care	<p>Inclusion:</p> <ul style="list-style-type: none"> ○ Patient requiring continuous albuterol therapy ○ CRS 0-5 at time of disposition and/or transfer, <i>if stable or improving on allowable max. therapies</i> <p>Exclusion:</p> <ul style="list-style-type: none"> ○ Patient <i>is not</i> stable or improving on allowable max. therapies ○ CPAP or BiPAP use for patients with <i>acute</i> respiratory disease

Goals and Outcome Measures

Process

- Rapid Response Team activation for reintubation

Outcome

- Therapy failure
- Length of stay

References

1. Arora, B., Mahajan, P., Zidan, M. A., & Sethuraman, U. (2012). Nasopharyngeal airway pressures in bronchiolitis patients treated with high-flow nasal cannula oxygen therapy. *Pediatric Emergency Care*, 28(11), 1179-1184.
2. Beggs, S., Wong, Z. H., Kaul, S., Ogden, K. J., & Walters, J. A. (2014). High-flow nasal cannula therapy for infants with bronchiolitis. *Cochrane Database of Systematic Reviews*, 1, CD009609.
3. Bressan, S., Balzani, M., Krauss, B., Pettenazzo, A., Zanconato, S., & Baraldi, E. (2013). High-flow nasal cannula oxygen for bronchiolitis in a pediatric ward: a pilot study. *European Journal of Pediatrics*, 172(12), 1649-1656.
4. Hilliard, T. N., Archer, N., Laura, H., Heraghty, J., Cottis, H., Mills, K., et al. (2012). Pilot study of vapotherm oxygen delivery in moderately severe bronchiolitis. *Archives of Disease in Childhood*, 97(2), 182-183.
5. Hough, J. L., Pham, T. M., & Schibler, A. (2014). Physiologic effect of high-flow nasal cannula in infants with bronchiolitis. *Pediatric Critical Care Medicine*, 15(5), e214-219.
6. McKiernan, C., Chua, L. C., Visintainer, P. F., & Allen, H. (2010). High flow nasal cannulae therapy in infants with bronchiolitis. *Journal of Pediatrics*, 156(4), 634-638.
7. Metge, P., Grimaldi, C., Hassid, S., Thomachot, L., Loundou, A., Martin, C., et al. (2014). Comparison of a high-flow humidified nasal cannula to nasal continuous positive airway pressure in children with acute bronchiolitis: Experience in a pediatric intensive care unit. *European Journal of Pediatrics*, 173(7), 953-958.
8. Milesi, C., Baleine, J., Matecki, S., Durand, S., Combes, C., Novais, A. R. et al. (2013). Is treatment with a high flow nasal cannula effective in acute viral bronchiolitis? A physiologic study. *Intensive Care Medicine*, 39(6), 1088-1094.
9. Rubin, S., Ghuman, A., Deakers, T., Khemani, R., Ross, P., & Newth, C. J. (2014). Effort of breathing in children receiving high-flow nasal cannula. *Pediatric Critical Care Medicine*, 15(1), 1-6.
10. Schibler, A., Pham, T. M., Dunster, K. R., Foster, K., Barlow, A., Gibbons, K., et al. (2011). Reduced intubation rates for infants after introduction of high-flow nasal prong oxygen delivery. *Intensive Care Medicine*, 37(5), 847-852.
11. Spentzas, T., Minarik, M., Patters, A. B., Vinson, B., & Stidham, G. (2009). Children with respiratory distress treated with high-flow nasal cannula. *Journal of Intensive Care Medicine*, 24(5), 323-328.
12. Wing, R., James, C., Maranda, L. S., & Armsby, C. C. (2012). Use of high-flow nasal cannula support in the emergency department reduces the need for intubation in pediatric acute respiratory insufficiency. *Pediatric Emergency Care*, 28(11), 1117-1123.
13. Abboud, P. A., Roth, P. J., Skiles, C. L., Stolfi, A., & Rowin, M. E. (2012). Predictors of failure in infants with viral bronchiolitis treated with high-flow, high-humidity nasal cannula therapy*. *Pediatric Critical Care Medicine*, 13(6), e343-349.
14. Kelly, G. S., Simon, H. K., & Sturm, J. J. (2013). High-flow nasal cannula use in children with respiratory distress in the emergency department: predicting the need for subsequent intubation. *Pediatric Emergency Care*, 29(8), 888-892.
15. Mayfield, S., Bogossian, F., O'Malley, L., & Schibler, A. (2014). High-flow nasal cannula oxygen therapy for infants with bronchiolitis: Pilot study. *Journal of Paediatrics and Child Health*, 50(5), 373-378.
16. Wraight, T. I., & Ganu, S. S. (2015). High-flow nasal cannula use in a paediatric intensive care unit over 3 years. *Critical Care & Resuscitation*, 17(3), 197-201.

Clinical Standards Preparation

This clinical standard was prepared by the Evidence-Based Outcomes Center (EBOC) team in collaboration with content experts at Texas Children's Hospital. Development of this clinical standard supports the TCH Quality and Patient Safety Program initiative to promote clinical standards and outcomes that build a culture of quality and safety within the organization.

HFNC Therapy Content Expert Team

Darlene Acorda, RN, CNS, Pediatric Intensive Care Unit
 Brian Bassham, MD, Emergency Center
 Dexter Buelow, RT
 Danny Castro, MD, Critical Care
 Jamie Causey, MD, Critical Care
 Charlene Davis, RN, CNS, Emergency Center
 Mindy Fein, MD, Emergency Center
 Bryan Greenfield, MD, Emergency Center
 Suzanne Iniguez, RT
 Jamie Jump, MD, Critical Care
 Sarah Meskill, MD, Emergency Center
 Brent Mothner, MD, Pediatric Hospital Medicine
 Vipul Parikh, MD, Pediatric Hospital Medicine
 Angela Stutts, RN, CNS, Pediatric Intensive Care Unit
 Cheryl Trumble-Wilkins, RN, CNS, Acute Care
 Jenny Werdenberg, MD, Pediatric Hospital Medicine
 Elizabeth Wuestner, RN, CNS, Emergency Center
 EBOC Team

Development Process

This clinical standard was developed using the process outlined in the EBOC Manual. The literature appraisal documents the following steps:

1. Review Preparation
 - PICO questions established
 - Evidence search confirmed with content experts
2. Review of Existing Internal and External Guidelines
 - N/A
3. Literature Review of Relevant Evidence
 - Searched: Cochrane, PubMed, Google
4. Critically Analyze the Evidence
 - 1 randomized controlled trial and 14 nonrandomized studies
5. Summarize the Evidence
 - Materials used in the development of the clinical standard, literature appraisal, and any order sets are maintained in a HFNC Therapy evidence-based review manual within EBOC.

Evaluating the Quality of the Evidence

Published clinical guidelines were evaluated for this review using the **AGREE II** criteria. The summary of these guidelines are included in the literature appraisal. AGREE II criteria evaluate Guideline Scope and Purpose, Stakeholder Involvement, Rigor of Development, Clarity and Presentation, Applicability, and Editorial Independence using a 4-point Likert scale. The higher the score, the more comprehensive the guideline. This clinical standard specifically summarizes the evidence *in support of* or *against* specific interventions and identifies where evidence is

lacking/inconclusive. The following categories describe how research findings provide support for treatment interventions.

"Evidence Supports" provides evidence to support an intervention

"Evidence Against" provides evidence against an intervention.

"Evidence Lacking/Inconclusive" indicates there is insufficient evidence to support or refute an intervention and no conclusion can be drawn *from the evidence*.

The **GRADE** criteria were utilized to evaluate the body of evidence used to make practice recommendations. The table below defines how the quality of the evidence is rated and how a strong versus weak recommendation is established. The literature appraisal reflects the critical points of evidence.

Recommendation	
STRONG	Desirable effects clearly outweigh undesirable effects or vice versa
WEAK	Desirable effects closely balanced with undesirable effects
Quality	Type of Evidence
High	Consistent evidence from well-performed RCTs or exceptionally strong evidence from unbiased observational studies
Moderate	Evidence from RCTs with important limitations (e.g., inconsistent results, methodological flaws, indirect evidence, or imprecise results) or unusually strong evidence from unbiased observational studies
Low	Evidence for at least 1 critical outcome from observational studies, RCTs with serious flaws or indirect evidence
Very Low	Evidence for at least 1 critical outcome from unsystematic clinical observations or very indirect evidence

Recommendations

Practice recommendations were directed by the existing evidence and consensus amongst the content experts. Patient and family preferences were included when possible. The Content Expert Team and EBOC team remain aware of the controversies in the initiation and escalation of HFNC therapy in children. When evidence is lacking, options in care are provided in the clinical standard and the accompanying order sets (if applicable).

Approval Process

Clinical standards are reviewed and approved by hospital committees as deemed appropriate for its intended use. Clinical standards are reviewed as necessary within EBOC at Texas Children's Hospital. Content Expert Teams are involved with every review and update.

Disclaimer

Practice recommendations are based upon the evidence available at the time the clinical standard was developed. Clinical standards (guidelines, summaries, or pathways) do not set out the standard of care and are not intended to be used to dictate a course of care. Each physician/practitioner must use his or her independent judgment in the management of any specific patient and is responsible, in consultation with the patient and/or the patient's family, to make the ultimate judgment regarding care

Version History

Date	Comments
Sep 2016	Originally completed
Jan 2018	Changed CRS cutoff for HFNC therapy, removed hypertonic saline, and added Woodlands Acute Care to the table on p. 2
Feb 2023	Pathway and algorithm updated